

Real or Interloper? The Redshift Likelihoods of Galaxies in the HUDF12

Nor Pirzkal¹, Barry Rothberg^{2,3},
Russell Ryan¹, Dan Coe¹, Sangeeta Malhotra⁴, James Rhoads⁴, Kai Noeske¹

ABSTRACT

In the absence of spectra, fitting template model spectra to observed photometric fluxes, known as Spectral Energy Distribution (SED) fitting, has become the workhorse for identifying high- z galaxies. In this paper, we present an analysis of the most recent and possibly most distant galaxies discovered in the Hubble Ultra Deep Field using a more robust method of redshift estimation based on Markov Chain Monte Carlo fitting (MCMC), rather than relying on the redshift of "best fit" models obtained using common χ^2 minimization techniques. The advantage of MCMC fitting is the ability to accurately estimate the probability density function of the redshift, as well as any other input model parameters, allowing us to derive accurate credible intervals by properly marginalizing over all other input model parameters. We apply our method to 13 recently identified sources and show that, despite claims based on χ^2 minimization, none of these sources can be securely ruled out as low redshift interlopers given the low signal-to-noise of currently available observations. We estimate that there is an average probability of 21% that these sources are low redshift interlopers.

Subject headings: Methods: statistical, Galaxies: high-redshift, Galaxies: photometry, Galaxies: stellar content

1. Introduction

Over the last 15 years the ever increasing pace at which new, more sensitive detectors and larger telescope apertures have come online has spurred a fast and furious race to detect the

¹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD21218, USA

²Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482, Potsdam, Germany

³George Mason University, Department of Physics & Astronomy, MS 3F3, 4400 University Drive, Fairfax, VA 22030, USA

⁴School of Earth And Space Exploration, Arizona State University, Tempe, AZ, 85287-1404, USA

most distant galaxies in the Universe. Emission features from HII regions (such as Lyman- α), have been a useful tool for detecting star-forming galaxies at $3 < z < 6$ (corresponding to rest-frame optical wavelengths) (Malhotra et al. 2001). However, at higher redshifts, corresponding to the first several hundred million years since the Big Bang, more and more matter was locked up in neutral hydrogen. At these early epochs the neutral intergalactic medium (IGM) attenuates significant amounts of light from galaxies, including Lyman- α , making detection of these systems more difficult. If Lyman- α is significantly diminished then one must rely on the detection of a continuum break to select candidate galaxies. Detecting galaxies at these redshifts places constraints on the epoch of re-ionization, thought to be at $z > 6$ (Fan et al. 2006; Komatsu et al. 2011). Detecting and measuring the properties of these galaxies is critical to understand what caused the epoch of re-ionization.

Detecting and measuring the properties of high- z galaxies is not easy. At rest-frame $\lambda < 1216\text{\AA}$ the flux from high redshift galaxies can be completely vanquished due to the Lyman- α forest absorption and when observed in a filter corresponding to this wavelength the galaxy “drops out.” At $\lambda > 1216\text{\AA}$ the flux is not attenuated and can be detected. Such breaks become more pronounced with redshift, for example at $z > 5.7$ a break corresponds to ~ 3.4 mags in color (e.g. Songaila & Cowie 2002; Hu et al. 2004). Detection at $z > 8$ is further complicated by the fact that the Lyman regime is redshifted into near-infrared wavelengths ($\lambda > 1\text{ }\mu\text{m}$) where bright atmospheric telluric skylines and strong telluric absorption features force ground-based observations to use specific and restrictive atmospheric windows. Large HST based spectroscopic surveys such as GRAPES and PEARS (Pirzkal et al. 2004) have demonstrated the power of low resolution spectroscopy to identify high redshift sources (Malhotra et al. 2005; Rhoads et al. 2009).

Recently, the race to find the earliest galaxies has relied more and more on the use of the near-infrared Wide Field Camera 3 (WFC3) detector ($1 < \lambda < 1.6\text{ }\mu\text{m}$) on the Hubble Space Telescope (*HST*), and in particular, broad-band imaging. Detecting possible $1216\text{\AA}$ breaks with *HST* photometry should be straightforward, assuming one collects a sufficient amount of photons. However, observations must first contend with the possibility of foreground interlopers. At $z \sim 6$ foreground interlopers can be as frequent as 45% for observations with signal-to-noise (S/N) < 5 (e.g. Dickinson et al. 2004; Stanway et al. 2008)). Without the direct detection of spectroscopic lines, redshift confirmations must rely on the application of SED fitting to measured photometric fluxes.

One of the first tools used to derive photometric redshifts was the SED fitting code **hyperz** which simply maximizes the likelihood by brute-force on a grid with standard SED templates (Bolzonella et al. 2000). As noted in Bolzonella et al. (2000): “Both the z_{phot} and the SED are obtained through **hyperz**, together with the best fit parameters (A_V , spectral type, metallicity and age). Because of the degeneracy between these parameters, the relevant information shall be the redshift and the rough SED type, in the sense that a given object

has a “blue” or “red” continuum at a given z , but no reliable information can be obtained about the other parameters from broad-band photometry alone.” Despite of this, **hyperz** and other similar χ^2 minimization SED fitting codes (e.g. Thompson et al. 2001; Papovich et al. 2001; Labbé et al. 2003; Schaerer & Pelló 2005; Mobasher et al. 2005; Pirzkal et al. 2007; McLure et al. 2009) have been used to extract more than just redshifts, including these degenerate properties. All of the latter is based on simply selecting the template with the smallest χ^2 .

This has led to a number of situations in which photometric redshifts and galaxy properties determined from χ^2 minimization SED fitting have produced questionable claims of high- z galaxies. Pelló et al. (2004) claimed the discovery of a $z = 10$ galaxy from a combination of deep J , H , K photometry obtained from the VLT, *HST* optical imaging, and J -band spectroscopy from ISAAC/VLT. While Pelló et al. (2004) claimed a detection of Lyman- α from low S/N ISAAC/VLT spectra, the basis for the claim relies primarily on SED fitting from the application of **hyperz** to stacked J , H , K imaging. Followup deep V -band ($V_{AB} = 27.8$ mag) imaging from the VLT (Lehnert et al. 2005) and deeper H -band imaging from Gemini/NIRI (Bremer et al. 2004) failed to detect fluxes predicted from the SED fitting. Lehnert et al. (2005) and a re-analysis of the spectra by Weatherley, et al. (2004) led to the conclusion that the object was a spurious detection, even though the fits from **hyperz** produced an excellent χ^2 fit. Mobasher et al. (2005) claimed a strong Balmer photometric break between the H -band and Spitzer IRAC1 ($3.6 \mu\text{m}$) resulting in a $z = 6.5$ claim for an early-type quiescent galaxy and raising serious questions regarding the viability of the Λ -CDM paradigm of building bigger galaxies from smaller building blocks. The redshift claim was based solely on a better χ^2 fit to Bruzual & Charlot (2003) models (CB03) in lieu of a dusty starburst model at $z = 2.5$ (the best fit from a secondary χ^2 minimum). When combined with followup Spitzer IRS data at 16 and $22 \mu\text{m}$, revised χ^2 minimization SED fitting produced a best fit for a $z = 1.7$ luminous infrared galaxy (Chary et al. 2007). Similarly, Henry et al. (2008) claimed the discovery of a Lyman break galaxy at $z \sim 9$ from NICMOS/*HST* imaging (based on a *F110W* dropout) and Spitzer imaging. Their χ^2 minimization fitting rejected local minima models at $z \sim 2$ -3 (produced by the degeneracy among the parameters). Subsequently, it was observations from MMT/Megacam blueward of the claimed break ($i' = 26$ mag) that demonstrated the intermediate redshift solutions that were initially rejected were actually correct (Henry et al. 2009). Finally, Capak et al. (2011) presented three z -band dropout candidates in which χ^2 minimization fitting produced best solutions at $z > 7$, only to find that followup imaging of one blueward of the claimed break and at $24 \mu\text{m}$ better supported at $z \sim 1.7$ solution; and a second with observed H -band flux (from NIRSPEC/Keck spectroscopy) weaker than predicted by SED models.

One should of course always keep in mind that any SED fitting effort is further complicated by the assumption that model galaxy spectra constructed from our knowledge of how

stars evolve in the local Universe is equally applicable to the earliest epochs. One should also be cognizant of the fact that the rest-frame wavelength range sampled by a fixed set of photometric filters will span a narrower range with increasing redshift. For the most recent claims of galaxies at $8 < z < 12$ (e.g. Yan et al. 2010; Bouwens et al. 2011; Ellis et al. 2013), WFC3/IR covers only $\Delta\lambda \sim 560$ and $390 \text{ \AA}$, (for $z = 8$ and $z = 12$, respectively, and excluding the $1216\text{\AA}$ break itself). Including the less sensitive and lower resolution IRAC1 and IRAC2 channels only widens this to $\Delta\lambda \sim 3700$ and $2600 \text{ \AA}$ for $z = 8$ and 12 , respectively. Strong statistical priors exist for only a handful of lensed galaxies (e.g. Coe et al. 2013). The relative lack of information about high redshift galaxy candidates, as well as the low signal-to-noise of the observations produce a situation where using sound statistical methods is particularly important when trying to determine the redshifts of these objects.

The latest claims of $z > 8$ galaxies based on χ^2 minimization SED fitting have been presented by Bouwens et al. (2011), Yan et al. (2010), and Ellis et al. (2013) using deep WFC3/IR images of the Hubble Deep Field. The results from Ellis et al. (2013) are particularly interesting since this project adds additional data (GO 12498, PI: Ellis, HUDF12) to the HUDF09 field (GO 11563, PI: Illingworth) in the F105W and F160W filters as well as new observations with the F140W filter. This filter spans a wavelength range inaccessible from the ground. Surprisingly since the fields are nearly identical, all but one of the sources listed in Bouwens et al. (2011) and none of the sources listed in Yan et al. (2010) were confirmed using these new deeper data. The new near-infrared measurements from the HUDF12 are the deepest observations available to date and for the foreseeable future. When combined with ancillary data from the HUDF09, these data allow one to construct spectral energy distributions of these sources spanning from observed B -band (ACS $F435W$) to IRAC1 and IRAC2 channels (3.4 and $4.6 \text{ }\mu\text{m}$, respectively). The 13 objects listed in Ellis et al. (2013) show large red colors consistent with strong spectral breaks. While followup spectroscopic observations have only been attempted on one of the targets, UDFy-38135539 (Lehnert et al. 2010; Bunker et al. 2013) and will be discussed later, the objects have all been fit using what is now considered the “standard method” of minimized χ^2 SED fitting, yielding redshifts of $8 < z < 12$. Given the limitations of this method, and the numerous examples of what later turned out to be incorrect redshift identifications, we are motivated to apply more robust Bayesian techniques to better constrain the physical parameters of galaxies observed using broad band filters. Central to this effort is the need to derive realistic estimates of the uncertainties for each derived parameter.

Thus, using the sources and fluxes, including ACS, WFC3 and IRAC, provided in Ellis et al. (2013), and reference therein, for 13 candidate high- z $1216\text{\AA}$ break galaxies we employ πMC^2 , a Markov Chain Monte Carlo (MCMC) approach (for details see Pirzkal et al. 2012a,b). For the purpose of this paper we define low redshift to be $z < 4$ and high redshift

to be $z > 4$. Further, we treat all 13 objects listed in Ellis et al. (2013) as potential high redshift candidates to be tested with πMC^2 . We are not implying that πMC^2 is the only viable method to perform this type of analysis. MCMC is a well established approach to this type of problem. The marginal signal-to-noise of current high redshift candidates observations warrant more robust techniques than the common χ^2 minimization technique and the reliance on a single “best fit” model provided by **hyper**-derived SED fitting software.

2. Applied Techniques: Minimized χ^2 SED fitting vs. πMC^2

Nominally, the “standard method” of minimized χ^2 SED fitting works by matching photometric fluxes to a pre-selected grid of model templates until the residuals between model and data are minimized. However, the choice of input models, the number of parameters (and whether any are constrained), parameter degeneracy and if the models sufficiently sample the entire range of the physical parameter space all affect the outcome of the fit. Any and all of these issues can produce “best-fit” solutions that are unphysical (i.e. a very old object in a young Universe as in Labbé et al. (2010); Richard et al. (2011)). Once the number of model parameters to be fit exceeds three it becomes computationally inefficient to fit parameter grid SEDs, more so for errors because each parameter must be refit following a pure Monte Carlo approach (e.g. each parameter requires several thousand additional iterations). All of the most recent *HST* based high- z candidate objects have relied on some modified implementation of the **hyperz** photometric code, or similar χ^2 SED minimization techniques (e.g. McLure et al. 2010, 2011; Bouwens et al. 2011; Ellis et al. 2013). While **hyperz** has proven to be capable for accurately constraining low- z photometric redshifts for large numbers of sources in deep, often crowded fields later confirmed with spectroscopic followup, it was designed to estimate the *most likely* photometric redshift, not produce credible intervals. Its treatment of the probability density functions (PDFs) is inadequate. Based on the **hyperz** Users Manual and Miralles (2013) it minimizes over the extra dimensions ($\mathbf{T}$) instead of marginalizing over them, therefore the PDF returned by **hyperz** is given by:

$$p_{\text{hyperz}}(z) = \min_{\mathbf{T}} p(z, \mathbf{T}) \quad (1)$$

This approach does not account for the *volume* of the parameter space probed resulting in unreliable confidence intervals, particularly the case of additional maxima. While such a distribution may be somewhat useful, the credible intervals produced by **hyperz** should not be trusted (Miralles 2013).

In contrast, MCMC, and our implementation of it, πMC^2 , is a random sampling method which samples the entire parameter space but does not sample the multidimensional region

uniformly. The posterior probability distribution function can be constructed by simply creating a histogram of the values of a given parameter in the MCMC chain from values taken after the chain converges (see Section 2 in Pirzkal et al. (2012a) for more details). Just as in Pirzkal et al. (2012a) we quote both the 95% and 68% credible regions here (see Section 2.1 of Pirzkal et al. (2012a) for a discussion of 95% vs 68% credible regions, which we take to be the 95% and 68% Highest Posterior Density, or HPD, regions, respectively). While computationally more expensive (yet more efficient), our current implementation of the MCMC SED fitting code, πMC^2 , has three main features:

- 1) It does not depend on a pre-defined input model parameter grid and allows for a computationally efficient exploration of the input parameter space.
- 2) the effect of nebular emission lines and nebular emission continuum can be included.
- 3) It allows one to derive a statistically valid posterior PDF for each of the input parameters by integrating over the remaining parameters:

$$p(z) = \int p(z, \mathbf{T}) d\mathbf{T} \quad (2)$$

To properly assess the possibility of lower-redshift solutions, it is imperative that the volume of parameter space be taken into account by integrating over the additional degrees of freedom.

We have applied πMC^2 to the sample of high redshift candidates listed in Ellis et al. (2013) because this sample is based on the deepest *HST* observations to date and provides candidate objects that could be the faintest examples of Lyman break galaxies. The analysis of redshifts and other parameters using the photometric fluxes from Ellis et al. (2013) are based upon using the use of stellar population models from Bruzual & Charlot (2003) (BC03), Charlot and Bruzual (2013) (CB07), and Maraston (2005) (MA05) in conjunction with πMC^2 . The parameters investigated include: stellar mass; a broad range of extinction values ($0 < A_v < 10$); metallicity ($0.005 < Z < 0.05$); and stellar population ages (limited only by the age of the Universe at a given redshift). We adopted flat priors for these parameters. The application of these models has also been tested with and without the effects of nebular continuum and emission lines (as parametrized by the escape fraction parameter). Details regarding the application of continuum nebular emission and emission lines to πMC^2 can be found in Section 2.2 of Pirzkal et al. (2012a). In addition to using single stellar population models (SSP), the BC03 models were also tested using an exponentially decaying star formation history model (parametrized by τ). Thus, there are seven discrete population models used with πMC^2 :

- 1) A SSP BC03 model with nebular contribution; 2) A SSP BC03 model without nebular contribution; 3) A BC03 model with $e^{-t/\tau}$ star-formation history and without nebular contribution; 4) A SSP CB07 model with nebular contribution; 5) A SSP CB07 model without

nebular contribution; 6) A SSP M05 model with nebular contribution; and 7) A SSP M05 model without nebular contribution.

3. Results

3.1. Redshift Constraints

A subset of the results of the πMC^2 analysis, obtained using the BC03 stellar population model with nebular emission, is presented in the first five columns of Table 1. For the most part, the redshift, extinction, stellar ages, stellar masses and metallicities ranges are uniform among the seven input models we used with πMC^2 . Any particular outliers are noted in Table 1.

The results using the BC03 are representative of what is derived for nearly all of the models. Table 1 lists the 68% and the 95% credible intervals derived using πMC^2 . Table 1 clearly shows that the 95% credible regions do *not* strongly constrain the redshift ranges of most of the sources. It is also clear from Table 1 that the 95% credible intervals are much larger than the 68% credible intervals which means that redshift posterior PDFs are *not* Gaussian.

A comparison of the quality of the fits (parameterized as the log likelihood computed by MCMC) from the most representative high redshift ($z > 4$) solution and that of the most representative low redshift ($z < 4$) solution, shows that high redshift models *appear* to fit the observations better than lower redshift models. As a first step, we can examine whether high redshift models always fit the data *significantly* better than alternative low redshift models by computing the significance p of a likelihood ratio test. This is a comparison between the log likelihood of the best fitting (usually $z > 4$) and best fitting low- z ($z < 4$) models. If $p > 0.05$ then the hypothesis that the high- z model fits the observation better than the low- z model is rejected at the 2σ level.

While we used stellar population models both with and without the effect of nebular lines is significant, we find that when using BC03 models without nebular emission, 7 of the 13 sources are reasonably fit by low redshift models. However, if nebular emission is included then *12 out of 13* objects are reasonably fit by low redshift models. The full range of values obtained from using different stellar population models are shown in Table 1. Clearly, the quality of the fits is not enough to confidently rule out these sources as low redshift interlopers.

The MCMC methodology also allows us to compute the *actual* probability that a source is at $z < 4$ by integrating the posterior PDF of each object. The range of values across all seven stellar population models described above is shown in the last column of Table 1. Low

redshift solutions *cannot* be strongly excluded based on the observed photometric break for many of the objects listed in Table 1. When examining our results using all seven stellar population models, We find that 9 out of 13 objects cannot be ruled out as low redshift interlopers in the best case scenario (i.e. selecting the stellar population models that most favor high redshift solutions), and none out of 13 in the worst case scenario (i.e. selecting the stellar population models that most favor low redshift solutions).

Low redshift ($z < 4$) solutions can, at best, be excluded at the 2σ confidence level ($P(z < 4) < 0.05$) for only four sources (UDF12-3947-8076, UDF12-3954-6284, UDFy-3779600 and UDFy-38135539). For each of the seven stellar population, the average probability for $P(z < 4)$, averaged across all 13 sources, is remarkably consistent and ranges from 19% to 26% with a global average of 21%. We conclude that $\approx 21\%$ of the sources in Table 1 are therefore likely to be low redshift interlopers.

As in the case of earlier attempts to detect high- z systems (e.g. Henry et al. 2009), it turns out that the limiting sensitivities of fluxes *blueward* of the presumptive break (i.e. ACS bands in our case) are not sufficient to provide strong constraints for the redshifts proposed by simple minimized χ^2 and "best fit" model SED fitting. Similarly, the IRAC observations are too shallow to provide enough constraints on the rest frame optical light from most of these objects.

In the case of four sources for which low- z solutions can be *statistically* rejected, three (UDF12-3947-8076, UDFy-3779600 and UDFy-38135539) are y-band dropouts and the brightest objects in the sample. Thus, relative to the ACS and IRAC detection limits, they have enough flux in the WFC3 bands to produce a change in flux strong enough to resemble a 1216Å decrement. The third object, UDF12-3954-6284, is very faint and only detected in *one* band. By definition it is a marginally acceptable candidate because it is only detected in one band (as noted by Ellis et al. 2013). It is possible that the observed photometric break could be caused by a strong emission line at low- z (see the Discussion section).

The ACS and IRAC detection limits are too high to unambiguously identify a 1216Å decrement for the remaining 9 objects in the sample. We estimate that the ACS and IRAC detection limits are ≈ 5 and ≈ 3 times too high (respectively) to distinguish between a 1216Å decrement and a Balmer break in sources this faint. This remains the main limiting factor in securely identifying sources at $z > 8.5$ using WFC3 observations of the HUDF.

In Figure 1 we show the SEDs of objects UDF12-3954-6284 and UDFy-37796000. The first example is meant to illustrate how too high of rest frame optical limits do not allow to reject low redshift solutions. The second example shows a clear detection of a strong photometric break as well as constraining limits in both the rest frame UV and optical bands. In the case of UDF12-3954-6284, the IRAC detection limits are clearly too high to rule out that this object is a low redshift interloper, if we allow for nebular emission. The log likelihood of the high- z and low- z solutions for this object are 720.9 and 722.4 with a likelihood

ratio test confidence of 0.08 which indicates that the high redshift model does not fit the observation significantly (i.e. 2σ) better than the low redshift model. While models lacking nebular emission clearly favor a high redshift solution for this object, the break can clearly be reproduced by an emission line.

A low redshift solution is much less likely in the case of the significantly brighter object UDFy-37796000 with a redshift of ≈ 1.6 and $A_V = 2.8$. The log likelihoods of the high and low redshift solutions are 722.6 and 717.5, respectively, resulting in a very low likelihood ratio test confidence values. This case demonstrates how strong rest-frame near UV and optical limits can help rule out a Balmer break.

3.2. Secondary Parameters

In addition to redshift constraints πMC^2 was used to determine other parameters including: stellar population ages; extinction; metallicity; and stellar mass. In the cases where models included nebular continuum, the escape fractions were determined; and in the case of BC03 with an exponentially decaying starburst, value of τ were derived. The 95% and 68% credible regions estimates for these parameters are given in Table 1 for each object. In the cases of three parameters: metallicity, escape fraction, and τ the posterior PDFs were essentially flat. This confirms results from Pirzkal et al. (2012a) where it was demonstrated that these parameters can only be constrained using very high precision photometry (i.e. better than 1% level), which is not the case for the UDF12.

The interdependence of the input model parameters is illustrated in Figure 2 which shows the two dimensional distribution of the posterior PDF's for UDF12-3895-7114. The statistically more likely regions are shown using proportionally lighter shades. This Figure illustrates the statistical complexity of fitting high redshift sources to stellar population models, non Gaussian nature of many of the input model parameter PDF's, as well as the low redshift solutions which fit the observations. As suggested above, deeper ACS observations (an increase in sensitivity by a factor of 5 for the *F850LP* ACS filter) would make it possible to more confidently exclude many of the low redshift solutions.

4. Discussion

The low signal-to-noise and large errors associated with the deepest WFC3 observations to date requires a more robust analysis than common χ^2 minimization techniques can provide to derive accurate credible regions for redshifts, especially in the faint target, highest

redshift regime. MCMC is able to provide just such an analysis. A complete SED analysis of the high redshift candidates discussed in this paper shows that there is a significant possibility (1 in 5) that these sources are low to moderate redshift ($z < 4$) interlopers. This is in contrast to the results presented by Ellis et al. (2013), where they conclude that 7 out of 13 sources can be confidently classified as high- z galaxies. Examining our 68% redshift credible regions might get the impression that our results agree well with those of Ellis et al. (2013). However, the limited number of measurements as well as the low signal to noise of the available measurements cause the redshift PDF of these sources to be highly non gaussian. This in turns lead us to revise upward the probability that these sources are low redshift interlopers. In the end, we find that only the redshifts of the brightest y-dropouts in this sample are likely to be at a high redshift ($z \approx 8$). Although the 128 orbits of the UDF12 have added considerable photometric depth to the UDF, the sources listed in Table 1 are still extremely faint. Moreover, given the large uncertainties in both flux and position, current IRAC observations do not strongly constrain the redshift of the vast majority of these objects. Similarly, the ACS observations are of insufficient depth to strongly constrain the SED blue-ward of the (presumptive) Lyman (or possibly Balmer) break. Spectroscopic observations, while difficult and time consuming, still remain a viable method for further constraining the proposed photometric redshifts and have been attempted for two of the sources in Table 1. The first, UDF12-3954-6284, was observed using the WFC3 G141 grism ($\lambda = 1.075\text{--}1.7 \mu\text{m}$ with $\delta\lambda = 0.00465 \mu\text{m}$) as part of the 3D-HST survey (GO-12177; PI: van Dokkum) for two visits (4.7 ks each, the same orientation for each visit) and as a part of the CANDELS supernova follow-up program (GO-12099; PI: Riess) also for two visits (6.6 ks and 15 ks) each with a different orientation. Brammer et al. (2013) report a 2.7σ detection of an emission line at $1.599\mu\text{m}$ based on smoothing and cross correlating the 2D spectrum with a kernel constructed from the central $0''.3$ of the *F160W* thumbnail image of UDF12-3954-6284 and removing any contamination using a contamination model (see their Figure 2). The object is not detected in any single visit. If the emission line were from Lyman- α it would place the galaxy at $z \sim 12$, but Brammer et al. (2013) reject this possibility based not on the rest-frame equivalent width ($\sim 170 \text{ \AA}$) but because Lyman- α emission should be attenuated early in the epoch of re-ionization. Instead, they conclude the line is most likely [OIII], placing the object at $z \sim 2.2$. Note that, UDF12-3954-6284, which was detected in only one WFC3 broad band filter, is one of the three objects with the strongest probabilities of being at high redshift.

The second object with follow-up spectroscopic observations is UDFy-38135539, all of which were obtained with ground-based instruments. It was first observed with SINFONI on the VLT (integration time of 14.8 hours), centered on the *J*-band ($1.1\text{--}1.4 \mu\text{m}$) resulting in a reported 6σ detection of Lyman- α , placing the galaxy at $z = 8.555$ (Lehnert et al. 2010). Subsequent observations using X-SHOOTER ($0.3\text{--}2.5 \mu\text{m}$) on the VLT and MOIRCS

(0.9-1.78 μm) on Subaru (5 hours and 11 hours integration times, respectively), as well as re-analysis of the original SINFONI have disputed this claim based on non-detections of Lyman- α (Bunker et al. 2013). They further noted that the *F105W* WFC3 photometry is inconsistent with both the line flux claimed by Lehnert et al. (2010) and the redshift from McLure et al. (2009). If the redshift were correct, then the *F105W* flux should be detected at 15σ . Evidently, the discrepancy in photometric measurements casts a doubt on our previous conclusion that this object was more unlikely to be a low redshift interloper.

Using πMC^2 we find that there are two populations of low- z galaxies which fit the available data as well as the high- z solutions. The first are low-moderate dusty systems at $z \sim 2$, the second are significantly dustier objects (up to $A_V = 4$) at $z < 1$. Distinguishing between these low- z solutions requires significantly more sensitive/deeper ACS and IRAC observations (see Figure 1) than are currently available (note: these would also serve to reject or improve the likelihood of high- z redshifts as well). Deeper IRAC observations, should any be undertaken in the remaining time of the Spitzer Warm Mission, must contend with reaching source confusion limits as well as the higher background noise from operating in “warm mode.” Contrary to common wisdom and the conclusions reached by other high- z studies, deeper observations blue-ward (e.g. *F850LP* or *F814W*) of the presumptive break would be extremely useful. (see Figure 1). Alternatively, one can seek sources that are brightened by lensing due to foreground galaxy clusters to get more flux from high- z sources (e.g. Coe et al. 2013). However, only a limited volume can be probed using this method.

Unfortunately, deeper observations in the optical bands using ACS are not likely considering the prohibitive observing times that would be required. Slitless spectroscopic observations using WFC3 with the G141 grism would provide an excellent method of rejecting or supporting the claims of high redshifts. However, this requires multiple position angles to avoid contamination from other objects or observational artifacts in the field as well as observations significantly deeper than those attempted to date (Pirzkal et al. 2012b). The G141 WFC3 grism observations would have the sensitivity to detect the emission lines potentially responsible for the observed photometric breaks. For example, in the case of object UDF12-3954-6284, an emission line with a flux as low as $\approx 1.5 \times 10^{-18}$ erg/s/Å/cm² could be solely responsible for a 29.3 magnitude detection in the F160W filter. Such a faint line would require upward of forty-two orbits to be *significantly* detected. In the absence of deeper HST or IRAC observations, observers will have to await the launch of JWST. We estimate that 10Ks observations at 0.9 μm using NIRCAM with the JWST would be deep enough to confirm that the observed photometric breaks are strong enough to be 1216Å breaks. Until such observations are made, the nature as well as the volume density of very high redshift candidates are likely to remain uncertain unless significantly deeper ACS or WFC3 observations are taken. At the very least, our analysis suggests that the uncertainties of current estimates of the volume density of these objects should be adjusted to account for

the possibility that nearly one in four of these objects is not a high redshift galaxy.

5. Acknowledgment

We would like to warmly thank J. M. Miralles for a fruitful discussion of the working of `hyperz`. We would also like to thank F. Pierfederici for his contributions that ultimately allowed us to run πMC^2 on distributed computers using Celery and HT-Condor. Finally, we thank Dr. Richard Ellis and Dr. Rychard Bouwens and their collaborators for constructive and insightful comments.

REFERENCES

- Beckwith, S. V. W., Stiavelli, M., Koekemoer, A. M., et al. 2006, *AJ*, 132, 1729
- Bolzonella, M., Miralles, J.-M., & Pelló, R. 2000, *A&A*, 363, 476
- Bouwens, R. J., Illingworth, G. D., Labbe, I., et al. 2011, *Nature*, 469, 504
- Brammer, G. B., van Dokkum, P. G., Illingworth, G. D., et al. 2013, *ApJ*, 765, L2
- Bremer, M. N., Jensen, J. B., Lehnert, M. D., et al. 2004, *ApJ*, 615, L1
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Bunker, A. J., Caruana, J., Wilkins, S. M. et al. 2013, *MNRAS* *in press*
- Capak, P., Mobasher, B., Scoville, N. Z., et al. 2011, *ApJ*, 730, 68
- Charlot, S., Bruzual, G., 2013, in prep
- Chary, R. R., Teplitz, H. I., Dickinson, M. E. et al. 2007, *ApJ*, 665, 257
- Coe, D., Zitrin, A., Carrasco, M., et al. 2013, *ApJ*, 762, 32
- Dickinson, M., Stern, D., Giavalisco, H. C. et al. 2004, *ApJ*, 600, 99
- Ellis, R. S., McLure, R. J., Dunlop, J. S., et al. 2013, *ApJ*, 763, L7
- Fan, X., Strauss, M. A., Becker, R. H., et al. 2006, *AJ*, 132, 117
- Henry, A. L., Malkan, M. A., Colbert, J. W., et al. 2008, *ApJ*, 680, L97
- Henry, A. L., Siana, B., Malkan, M. A. et al. 2009, *ApJ*, 697, 1128
- Hu, E. M., Cowie, L. L., Capak, P. et al. 2004, *AJ*, 127, 563




Fig. 1.— High and low redshift solutions to the observed photometry of objects UDF12-3954-6284 and UDFy-37796000. The upper left sub-panels show all of the available photometry, including the infrared IRAC upper limits. The main panels show the observed UV to NIR (ACS and WFC3 bands) region for the same object. Two models are shown in each case. The first, shown in a dashed red line, is a $z > 8$ model while the solid blue line shows a much lower redshift solution at $z \approx 2$. Observed fluxes are shown using error bars. Synthetic model fluxes in each observed filter are shown using large circles.




Fig. 2.— Posterior probability density function estimates for object UDF12-3895-7114. The non-gaussian, multi-modal nature of the redshift distribution is evident in the rightmost plots.

Komatsu, E., Smith, K. M., Dunkley, J., et al. 2011, ApJS, 192, 18

Labbé, I, Franx, M., Rudnick, G. et al. 2003, AJ, 125, 1107

Labbé, I, Gonzalez, V., Bouwens, R. J. et al. 2010, ApJ, 716, 103

Lehnert, M. D., Förster-Schreiber, N. M., & Bremer, M. N. 2005, ApJ, 624, 80

Lehnert, M. D., Nesvadba, N. P. H., Cuby, J. G., et al. 2010, Nature, 467, 940

Maraston, C. 2005, MNRAS, 362, 799

Malhotra, S., Rhoads, J., Dey, A., Stern, D., & Spinrad, H. 2001, Gas and Galaxy Evolution, 240, 97

Malhotra, S., Rhoads, J. E., Pirzkal, N., et al. 2005, ApJ, 626, 666

Miralles, J-M. 2013, private communication.

McLure, R. J., Cirasuolo, M., Dunlop, J. S. et al. 2009, MNRAS, 395, 2196

McLure, R. J., Dunlop, J. S., Cirasuolo, M., et al. 2010, MNRAS, 403, 960

McLure, R. J., Dunlop, J. S., de Ravel, L., et al. 2011, MNRAS, 418, 2074

Mobasher, B., Dickinson, M., Ferguson, H. C. et al. 2005, ApJ, 635, 832

Papovich, C., Dickinson, M., Ferguson, H. C. et al. 2001, ApJ, 559, 620

Pelló, R., Schaerer, D., Richard, J., et al. 2004, Å, 416, L35

Pirzkal, N., Malhotra, S., Rhoads, J. E., & Xu, C. 2007, ApJ, 667, 49

Pirzkal, N., Rothberg, B., Nilsson, K. K., et al. 2012, ApJ, 748, 122

Pirzkal, N., Rothberg, B., Chun, L., et al. 2012, arXiv:1208.5535, ApJ*submitted*

Pirzkal, N., Xu, C., Malhotra, S., et al. 2004, ApJS, 154, 501

Richard, J., Kneib, J. P., Ebeling, H. et al. 2011, MNRAS, 414, L31

Rhoads, J. E., Malhotra, S., Pirzkal, N., et al. 2009, ApJ, 697, 942

Schaerer, D., & Pelló, R. 2005, MNRAS, 362, 1054

Songaila, A., & Cowie, L. L. 2002, AJ, 132, 2182

- Stanway, E. R., Bremer, M. N., & Lehnert, M. D. 2008, MNRAS, 385, 493
- Thompson, R. I., Weymann, R. J., & Storrie-Lombardi, L. J. 2001, ApJ, 546, 694
- Weatherley, S. J., Warren, S. J., & Babbedge, T. S. R. 2004, *a*, 428, L29
- Yan, H.-J., Windhorst, R. A., Hathi, N. P., et al. 2010, Research in Astronomy and Astrophysics, 10, 867